


Patent Claims

1. Pyridinylanilides of the formula (I)



5 in which

R represents hydrogen, fluorine, chlorine, methyl or trifluoromethyl;

R¹, R² and R³ independently of one another each represents hydrogen, halogen, cyano, nitro,
amino, hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl;

10 or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl,
alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkyl-
sulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;

or represents in each case straight-chain or branched alkenyl or alkenyloxy having in
each case 2 to 6 carbon atoms;

15 or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy,
halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each
case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;

or represents in each case straight-chain or branched halogenoalkenyl or halogeno-
alkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different
20 halogen atoms;

or represents in each case straight-chain or branched alkylamino, dialkylamino, alkyl-
carbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocar-
bonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms
in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to
25 6 carbon atoms in the respective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;

or represents in each case the grouping $-C(Q^1)=N-Q^2$, wherein

Q¹ represents hydrogen, hydroxyl or C₁-C₄-alkyl, C₁-C₄-halogenoalkyl having 1
to 9 identical or different halogen atoms, or C₃-C₆-cycloalkyl and

30 Q² represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents C₁-
C₄-alkyl or C₁-C₄-alkoxy, each of which is optionally substituted by halogen,
cyano, hydroxyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino, di(C₁-C₄-

alkyl)amino or phenyl; or represents C₂-C₄-alkenyloxy or C₂-C₄-alkynyloxy, represents phenyl, phenoxy, phenylthio, benzoyl, benzoylethenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched C₁-C₄-alkyl and C₁-C₄-alkoxy;

or

R² and R³, if attached to the pyridinyl moiety in ortho position to each other, furthermore together represent C₃-C₄-alkylene, C₃-C₄-alkenylene, C₂-C₃-oxyalkylene or C₁-C₂-dioxyalkylene, in each case optionally mono- to tetra-substituted, identically or differently, by fluorine, chlorine, oxo, methyl, ethyl, trifluoromethyl;

R⁴ represents hydrogen, C₁-C₈-alkyl, C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₃-C₈-cycloalkyl; C₁-C₆-halogenoalkyl, C₁-C₄-halogenoalkylthio, C₁-C₄-halogenoalkylsulfinyl, C₁-C₄-halogenoalkylsulfonyl, halogeno-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₃-C₈-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms; formyl-C₁-C₃-alkyl, (C₁-C₃-alkyl)carbonyl-C₁-C₃-alkyl, (C₁-C₃-alkoxy)carbonyl-C₁-C₃-alkyl; (C₁-C₃-halogenoalkyl)carbonyl-C₁-C₃-alkyl, (C₁-C₃-halogenoalkoxy)carbonyl-C₁-C₃-alkyl having in each case 1 to 7 fluorine-, chlorine- and/or bromine atoms, (C₁-C₃-alkyl)carbonyl-C₁-C₃-halogenoalkyl, (C₁-C₃-alkoxy)carbonyl-C₁-C₃-halogenoalkyl having in each case 1 to 6 fluorine-, chlorine- and/or bromine atoms, (C₁-C₃-halogenoalkyl)carbonyl-C₁-C₃-halogenoalkyl, (C₁-C₃-halogenoalkoxy)carbonyl-C₁-C₃-halogenoalkyl having in each case 1 to 13 fluorine-, chlorine- and/or bromine atoms; -COR⁵, -CONR⁶R⁷ or -CH₂NR⁸R⁹,

R⁵ represents hydrogen, C₁-C₈-alkyl, C₁-C₈-alkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₃-C₈-cycloalkyl; C₁-C₆-halogenoalkyl, C₁-C₆-halogenoalkoxy, halogeno-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₃-C₈-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms; or -COR¹⁰,

R⁶ and R⁷ independently of one another each represent hydrogen, C₁-C₈-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₃-C₈-cycloalkyl; C₁-C₈-halogenoalkyl, halogeno-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₃-C₈-halogenocycloalkyl having in each case 1 bis 9 fluorine-, chlorine- and/or bromine atoms,

R⁶ and R⁷ furthermore together with the nitrogen atom to which they are attached, represent a saturated 5- to 8-membered heterocycle, which heterocycle may have 1 or 2 additional, non-adjacent heteroatoms selected from the group consisting of oxygen, sulphur and NR¹¹, and which heterocycle may optionally be mono- to poly-substitu-

ted, identically or differently, by halogen or C₁-C₄-alkyl,

R⁸ and R⁹ independently of one another each represent hydrogen, C₁-C₈-alkyl, C₃-C₈-cycloalkyl; C₁-C₈-halogenoalkyl, C₃-C₈-halogenocycloalkyl having in each case 1 bis 9 fluorine-, chlorine- and/or bromine atoms,


5 R⁸ and R⁹ furthermore together with the nitrogen atom to which they are attached, represent a saturated 5- to 8-membered heterocycle, which heterocycle may have 1 or 2 additional, non-adjacent heteroatoms selected from the group consisting of oxygen, sulphur and NR¹¹, and which heterocycle may optionally be mono- to poly-substituted, identically or differently, by halogen or C₁-C₄-alkyl,

10 R¹⁰ represents hydrogen, C₁-C₈-alkyl, C₁-C₈-alkoxy, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₃-C₈-cycloalkyl; C₁-C₆-halogenoalkyl, C₁-C₆-halogenoalkoxy, halogeno-C₁-C₄-alkoxy-C₁-C₄-alkyl, C₃-C₈-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms,

R¹¹ represents hydrogen or C₁-C₆-alkyl,

15

A represents a radical of the formula (A1)



(A1), wherein

R¹² represents hydrogen, cyano, halogen, nitro, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₃-C₆-cycloalkyl, C₁-C₄-halogenoalkyl, C₁-C₄-halogenoalkoxy or C₁-C₄-halogenoalkylthio each having 1 to 5 halogen atoms, aminocarbonyl or aminocarbonyl-C₁-C₄-alkyl and

20


R¹³ represents hydrogen, halogen, cyano, C₁-C₄-alkyl, C₁-C₄-alkoxy or C₁-C₄-alkylthio and

R¹⁴ represents hydrogen, C₁-C₄-alkyl, hydroxy-C₁-C₄-alkyl, C₂-C₆-alkenyl, C₃-C₆-cycloalkyl, C₁-C₄-alkylthio-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-halogenoalkyl, C₁-C₄-halogenoalkylthio-C₁-C₄-alkyl, C₁-C₄-halogenoalkoxy-C₁-C₄-alkyl each having 1 to 5 halogen atoms, or phenyl,

25

or

A represents a radical of the formula (A2)



(A2), wherein

30


R¹⁵ and R¹⁶ independently of one another each represent hydrogen, halogen, C₁-C₄-

alkyl or C₁-C₄-halogenoalkyl having 1 to 5 halogen atoms and

R¹⁷ represents halogen, cyano or C₁-C₄-alkyl, or C₁-C₄-halogenoalkyl or C₁-C₄-halogenoalkoxy each having 1 to 5 halogen atoms,

or

5 A represents a radical of the formula (A3)



R¹⁸ and R¹⁹ independently of one another each represent hydrogen, halogen, C₁-C₄-alkyl or C₁-C₄-halogenoalkyl having 1 to 5 halogen atoms and

R²⁰ represents hydrogen, halogen, C₁-C₄-alkyl or C₁-C₄-halogenoalkyl having 1 to 5 halogen atoms,

or


A represents a radical of the formula (A4)



R²¹ represents hydrogen, halogen, hydroxyl, cyano, C₁-C₆-alkyl, C₁-C₄-halogenoalkyl, C₁-C₄-halogenoalkoxy or C₁-C₄-halogenoalkylthio each having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A5)



R²² represents halogen, hydroxyl, cyano, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-halogenoalkyl, C₁-C₄-halogenoalkylthio or C₁-C₄-halogenoalkoxy each having 1 to 5 halogen atoms and

R²³ represents hydrogen, halogen, cyano, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-halogenoalkyl, C₁-C₄-halogenoalkoxy each having 1 to 5 halogen atoms, C₁-C₄-alkylsulphinyl or C₁-C₄-alkylsulphonyl,

or


A represents a radical of the formula (A6)



- R^{24} represents C_1 - C_4 -alkyl or C_1 - C_4 -halogenoalkyl having 1 to 5 halogen atoms and
 R^{25} represents C_1 - C_4 -alkyl,
 Q^3 represents a sulphur or oxygen atom, represents SO , SO_2 or CH_2 ,
 p represents 0, 1 or 2, where R^{25} represents identical or different radicals if p represents 2,

or

A represents a radical of the formula (A7)




(A7), wherein

R^{26} represents C_1 - C_4 -alkyl or C_1 - C_4 -halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A8)




(A8), wherein

R^{27} represents C_1 - C_4 -alkyl or C_1 - C_4 -halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A9)




(A9), wherein

R^{28} and R^{29} independently of one another each represent hydrogen, halogen, amino, C_1 - C_4 -alkyl or C_1 - C_4 -halogenoalkyl having 1 to 5 halogen atoms and

R^{30} represents hydrogen, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A10)




(A10), wherein

R^{31} and R^{32} independently of one another each represent hydrogen, halogen, amino, nitro, C_1 - C_4 -alkyl or C_1 - C_4 -halogenoalkyl having 1 to 5 halogen atoms and

R^{33} represents hydrogen, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A11)




(A11), wherein

R^{34} represents hydrogen, halogen, amino, C_1 - C_4 -alkylamino, di- $(C_1$ - C_4 -alkyl)-amino, cyano, C_1 - C_4 -alkyl or C_1 - C_4 -halogenoalkyl having 1 to 5 halogen atoms and

R^{35} represents halogen, C_1 - C_4 -alkyl or C_1 - C_4 -halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A12)



(A12), wherein

R^{36} represents hydrogen, halogen, amino, C_1 - C_4 -alkylamino, di- $(C_1$ - C_4 -alkyl)-amino, cyano, C_1 - C_4 -alkyl or C_1 - C_4 -halogenoalkyl having 1 to 5 halogen atoms and

R^{37} represents halogen, C_1 - C_4 -alkyl or C_1 - C_4 -halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A13)




(A13), wherein

R^{38} represents halogen, C_1 - C_4 -alkyl or C_1 - C_4 -halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A14)




(A14), wherein

R^{39} represents hydrogen or C_1 - C_4 -alkyl and

R^{40} represents halogen or C_1 - C_4 -alkyl,

or

A represents a radical of the formula (A15)




(A15), wherein

R⁴¹ represents C₁-C₄-alkyl or C₁-C₄-halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A16)




(A16), wherein

R⁴² represents hydrogen, halogen, C₁-C₄-alkyl or C₁-C₄-halogenoalkyl having 1 to 5 halogen atoms,

or

A represents a radical of the formula (A17)



(A17), wherein

R⁴³ represents halogen, hydroxyl, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-halogenoalkyl, C₁-C₄-halogenoalkylthio or C₁-C₄-halogenoalkoxy each having 1 to 5 halogen atoms,

15 excluded compounds of the formula (I), in which


R represents hydrogen and

R¹, R² and R³ independently of one another each represents hydrogen, halogen; or straight-chain or branched alkyl having 1 to 4 carbon atoms; or straight-chain or branched halogenoalkyl having 1 to 4 carbon atoms; and

20 R⁴ represents hydrogen

and

A represents a radical of the formula (A1)



(A1), wherein


R¹² represents halogen, C₁-C₄-alkyl, C₁-C₄-halogenoalkyl and

R¹³ represents hydrogen and

R¹⁴ represents methyl,

or

A represents a radical of the formula (A2)




(A2), wherein

R¹⁵ and R¹⁶ independently of one another each represent hydrogen or C₁-C₄-alkyl and
R¹⁷ represents halogen, C₁-C₄-alkyl or C₁-C₄-halogenoalkyl,

5 or

A represents a radical of the formula (A4)




(A4), wherein

R²¹ represents halogen, C₁-C₄-alkyl or C₁-C₄-halogenoalkyl,

or

10

A represents a radical of the formula (A5)



(A5), wherein


R²² represents halogen and

R²³ represents hydrogen,

or

15

A represents a radical of the formula (A6)



(A6), wherein

R²⁴ represents methyl and


Q³ represents a sulphur or CH₂,

p represents 0,

20

or

A represents a radical of the formula (A9)



(A9), wherein


R²⁸ and R²⁹ independently of one another each represent hydrogen or C₁-C₄-alkyl and

R³⁰ represents methyl,

25

or

A represents a radical of the formula (A11)




(A11), wherein

R^{34} represents hydrogen or C_1 - C_4 -alkyl and

R^{35} represents halogen, C_1 - C_4 -alkyl or C_1 - C_4 -halogenoalkyl,

or

5 A represents a radical of the formula (A16)



(A16), wherein

R^{42} represents halogen.

2. Pyridinylanilides of the formula (I) according to Claim 1, in which

10 R represents hydrogen, fluorine, chlorine, methyl or trifluoromethyl;

R^1 , R^2 and R^3 independently of one another each represents hydrogen, halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl;

15 or represents in each case straight-chain or branched alkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 6 carbon atoms;

or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 4 carbon atoms and 1 to 9 identical or different halogen atoms;

20 or represents in each case straight-chain or branched alkylamino, dialkylamino, alkyl-carbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 4 carbon atoms in the respective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;

25 or represents the grouping $-C(Q^1)=N-Q^2$, wherein

Q^1 represents hydrogen, hydroxyl or C_1 - C_4 -alkyl, C_1 - C_4 -halogenoalkyl having 1 to 9 identical or different halogen atoms or C_3 - C_6 -cycloalkyl and

Q^2 represents hydroxyl, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -halogenoalkyl or C_1 - C_4 -halogenoalkoxy each having 1 to 9 identical or different halogen atoms,

30 or

R^2 and R^3 , if attached to the pyridinyl moiety in ortho position to each other, furthermore together represent $-(CH_2)_3-$, $-(CH_2)_4-$, $-CH=CH-CH=CH-$, $-O(CH_2)_2-$, $-O(CH_2)_3-$,


-OCH₂O-, -O(CH₂)₂O-, in each case optionally mono- to tetra-substituted, identically or differently, by fluorine, chlorine, oxo, methyl, ethyl, trifluoromethyl,

- 5 R⁴ represents hydrogen; C₁-C₆-alkyl, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl, C₁-C₃-alkoxy-C₁-C₃-alkyl, C₃-C₆-cycloalkyl; C₁-C₄-halogenoalkyl, C₁-C₄-halogenoalkylthio, C₁-C₄-halogenoalkylsulfinyl, C₁-C₄-halogenoalkylsulfonyl, halogeno-C₁-C₃-alkoxy-C₁-C₃-alkyl, C₃-C₆-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms; formyl-C₁-C₃-alkyl, (C₁-C₃-alkyl)carbonyl-C₁-C₃-alkyl, (C₁-C₃-alkoxy)carbonyl-C₁-C₃-alkyl; (C₁-C₃-halogenoalkyl)carbonyl-C₁-C₃-alkyl, (C₁-C₃-halogenoalkoxy)carbonyl-C₁-C₃-alkyl having in each case 1 to 7 fluorine-, chlorine- and/or bromine atoms, (C₁-C₃-alkyl)carbonyl-C₁-C₃-halogenoalkyl, (C₁-C₃-alkoxy)carbonyl-C₁-C₃-halogenoalkyl having in each case 1 to 6 fluorine-, chlorine- and/or bromine atoms, (C₁-C₃-halogenoalkyl)carbonyl-C₁-C₃-halogenoalkyl, (C₁-C₃-halogenoalkoxy)carbonyl-C₁-C₃-halogenoalkyl having in each case 1 to 13 fluorine-, chlorine- and/or bromine atoms; -COR⁵, -CONR⁶R⁷ or -CH₂NR⁸R⁹,
- 10 R⁵ represents hydrogen, C₁-C₆-alkyl, C₁-C₄-alkoxy, C₁-C₃-alkoxy-C₁-C₃-alkyl, C₃-C₆-cycloalkyl; C₁-C₄-halogenoalkyl, C₁-C₄-halogenoalkoxy, halogeno-C₁-C₃-alkoxy-C₁-C₃-alkyl, C₃-C₆-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms; or -COR¹⁰,
- 15 R⁶ and R⁷ independently of one another each represent hydrogen, C₁-C₆-alkyl, C₁-C₃-alkoxy-C₁-C₃-alkyl, C₃-C₆-cycloalkyl; C₁-C₄-halogenoalkyl, halogeno-C₁-C₃-alkoxy-C₁-C₃-alkyl, C₃-C₆-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms,
- 20 R⁶ and R⁷ furthermore together with the nitrogen atom to which they are attached, represent a saturated 5- to 8-membered heterocycle, which heterocycle may have 1 or 2 additional, non-adjacent heteroatoms selected from the group consisting of oxygen, sulphur and NR¹¹, and which heterocycle may optionally be mono- to tetra-substituted, identically or differently, by halogen or C₁-C₄-alkyl,
- 25 R⁸ and R⁹ independently of one another each represent hydrogen, C₁-C₆-alkyl, C₃-C₆-cycloalkyl; C₁-C₄-halogenoalkyl, C₃-C₆-halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms,
- 30 R⁸ and R⁹ furthermore together with the nitrogen atom to which they are attached, represent a saturated 5- to 8-membered heterocycle, which heterocycle may have 1 or 2 additional, non-adjacent heteroatoms selected from the group consisting of oxygen, sulphur and NR¹¹, and which heterocycle may optionally be mono- to tetra-substituted, identically or differently, by halogen or C₁-C₄-alkyl,
- 35

R^{10} represents hydrogen, C_1 - C_6 -alkyl, C_1 - C_4 -alkoxy, C_1 - C_3 -alkoxy- C_1 - C_3 -alkyl, C_3 - C_6 -cycloalkyl; C_1 - C_4 -halogenoalkyl, C_1 - C_4 -halogenoalkoxy, halogeno- C_1 - C_3 -alkoxy- C_1 - C_3 -alkyl, C_3 - C_6 -halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms,

5 R^{11} represents hydrogen or C_1 - C_4 -alkyl,

A represents a radical of the formula (A1)



(A1), wherein

10 R^{12} represents hydrogen, cyano, fluorine, chlorine, bromine, iodine, methyl, ethyl, iso-propyl, methoxy, ethoxy, methylthio, ethylthio, cyclopropyl, C_1 - C_2 -halogenoalkyl, C_1 - C_2 -halogenoalkoxy each having 1 to 5 fluorine, chlorine and/or bromine atoms, trifluoromethylthio, difluoromethylthio, aminocarbonyl, aminocarbonylmethyl or aminocarbonylethyl and

15 R^{13} represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, ethyl, methoxy, ethoxy, methylthio or ethylthio and

R^{14} represents hydrogen, methyl, ethyl, n-propyl, iso-propyl, C_1 - C_2 -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms, hydroxymethyl, hydroxyethyl, cyclopropyl, cyclopentyl, cyclohexyl or phenyl,

or

20 A represents a radical of the formula (A2)




(A2), wherein

R^{15} and R^{16} independently of one another each represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl or C_1 - C_2 -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms and

25 R^{17} represents fluorine, chlorine, bromine, cyano, methyl, ethyl, C_1 - C_2 -halogenoalkyl or C_1 - C_2 -halogenoalkoxy each having 1 to 5 fluorine, chlorine and/or bromine atoms,

or

A represents a radical of the formula (A3)



R^{18} and R^{19} independently of one another each represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl or C_1 - C_2 -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms and

5 R^{20} represents hydrogen, fluorine, chlorine, bromine, methyl, ethyl or C_1 - C_2 -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms,

or

A represents a radical of the formula (A4)



10 R^{21} represents hydrogen, fluorine, chlorine, bromine, iodine, hydroxyl, cyano, C_1 - C_4 -alkyl, C_1 - C_2 -halogenoalkyl, C_1 - C_2 -halogenoalkoxy or C_1 - C_2 -halogenoalkylthio each having 1 to 5 fluorine, chlorine and/or bromine atoms,

or

A represents a radical of the formula (A5)



15 R^{22} represents fluorine, chlorine, bromine, iodine, hydroxyl, cyano, C_1 - C_4 -alkyl, methoxy, ethoxy, methylthio, ethylthio, difluoromethylthio, trifluoromethylthio, C_1 - C_2 -halogenoalkyl or C_1 - C_2 -halogenoalkoxy each having 1 to 5 fluorine, chlorine and/or bromine atoms and

20 R^{23} represents hydrogen, fluorine, chlorine, bromine, iodine, cyano, C_1 - C_4 -alkyl, methoxy, ethoxy, methylthio, ethylthio, C_1 - C_2 -halogenoalkyl or C_1 - C_2 -halogenoalkoxy each having 1 to 5 fluorine, chlorine and/or bromine atoms, C_1 - C_2 -alkylsulphinyl or C_1 - C_2 -alkylsulphonyl,

or

25 A represents a radical of the formula (A6)




R^{24} represents methyl, ethyl or C_1 - C_2 -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms and

- R^{25} represents methyl or ethyl,
 Q^3 represents a sulphur atom, SO_2 or CH_2 ,
 p represents 0 or 1,

or

- 5 A represents a radical of the formula (A9)



R^{28} and R^{29} independently of one another each represent hydrogen, fluorine, chlorine, bromine, amino, methyl, ethyl or C_1 - C_2 -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms and

- 10 R^{30} represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, ethyl or C_1 - C_2 -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms,

or

- A represents a radical of the formula (A10)



- 15 R^{31} and R^{32} independently of one another each preferably represent hydrogen, fluorine, chlorine, bromine, amino, nitro, methyl, ethyl or C_1 - C_2 -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms and

R^{33} represents hydrogen, fluorine, chlorine, bromine, methyl, ethyl or C_1 - C_2 -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms,

20

or

- A represents a radical of the formula (A11)




R^{34} represents hydrogen, fluorine, chlorine, bromine, amino, C_1 - C_4 -alkylamino, di(C_1 - C_4 -alkyl)amino, cyano, methyl, ethyl or C_1 - C_2 -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms and

25

R^{35} represents fluorine, chlorine, bromine, methyl, ethyl or C_1 - C_2 -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms,

or

- A represents a radical of the formula (A12)




(A12), wherein

R^{36} represents hydrogen, fluorine, chlorine, bromine, amino, C_1 - C_4 -alkylamino, di(C_1 - C_4 -alkyl)amino, cyano, methyl, ethyl or C_1 - C_2 -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms and

5 R^{37} represents fluorine, chlorine, bromine, methyl, ethyl or C_1 - C_2 -halogenoalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms,

or

A represents a radical of the formula (A17)



(A17), wherein

10 R^{43} preferably represents fluorine, chlorine, bromine, iodine, hydroxyl, C_1 - C_4 -alkyl, methoxy, ethoxy, methylthio, ethylthio, difluoromethylthio, trifluoromethylthio, C_1 - C_2 -halogenoalkyl or C_1 - C_2 -halogenoalkoxy each having 1 to 5 fluorine, chlorine and/or bromine atoms,


15 excluded compounds of the formula (I), in which

R represents hydrogen and

R^1 , R^2 and R^3 independently of one another each represents hydrogen, halogen; or straight-chain or branched alkyl having 1 to 4 carbon atoms; or straight-chain or branched halogenoalkyl having 1 to 4 carbon atoms; and

20 R^4 represents hydrogen and

A represents a radical of the formula (A1)



(A1), wherein


25 R^{12} represents fluorine, chlorine, bromine, iodine, methyl, ethyl, iso-propyl, C_1 - C_2 -halogenoalkyl and

R^{13} represents hydrogen and

R^{14} represents methyl,

or

A represents a radical of the formula (A2)




(A2), wherein

R^{15} and R^{16} independently of one another each represent hydrogen, methyl or ethyl
and

R^{17} represents fluorine, chlorine, bromine, methyl, ethyl, or C_1 - C_2 -halogenoalkyl,

5 or

A represents a radical of the formula (A4)




(A4), wherein

R^{21} represents fluorine, chlorine, bromine, iodine, C_1 - C_4 -alkyl or C_1 - C_2 -halogenoalkyl,

10 or

A represents a radical of the formula (A5)




(A5), wherein

R^{22} represents fluorine, chlorine, bromine, iodine and
 R^{23} represents hydrogen,

15 or

A represents a radical of the formula (A6)



(A6), wherein

R^{24} represents methyl and


Q^3 represents a sulphur or CH_2 ,

20

P represents O ,

or

A represents a radical of the formula (A9)



(A9), wherein


R^{28} and R^{29} independently of one another each represent hydrogen, methyl or ethyl
and

R^{30} represents methyl,

25

or

A represents a radical of the formula (A11)



(A11), wherein

R^{34} represents hydrogen, methyl or ethyl and

5 R^{35} represents fluorine, chlorine, bromine, methyl, ethyl or C_1 - C_2 -halogenoalkyl.

3. Pyridinylanilides of the formula (I) according to Claim 1, in which

R represents hydrogen, fluorine, chlorine, methyl or trifluoromethyl;

10 R^1 , R^2 and R^3 independently of one another each represents hydrogen, fluorine, chlorine, bromine, cyano; methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl, methoxy, ethoxy, n- or isopropoxy, n-, iso-, sec- or tert-butoxy, methylthio, ethylthio, n- or iso-propylthio, n-, iso-, sec- or tert-butylthio, trifluoromethyl, trifluoroethyl, difluoromethoxy, trifluoromethoxy, difluorochloromethoxy, trifluoroethoxy, cyclopropyl, cyclopentyl, cyclohexyl,

15 or represents the grouping $-C(Q^1)=N-Q^2$, wherein

Q^1 represents hydrogen, methyl, ethyl, trifluoromethyl or cyclopropyl, and

Q^2 represents hydroxyl, methoxy, ethoxy, n-propoxy or iso-propoxy,

or

20 R^2 and R^3 , if attached to the pyridinyl moiety in ortho position to each other, furthermore together represent $-(CH_2)_3-$, $-(CH_2)_4-$, $-CH=CH-CH=CH-$, $-OCH_2O-$, $-O(CH_2)_2O-$, $-OCF_2O-$, $-O(CF_2)_2O-$,

25 R^4 represents hydrogen, methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl, pentyl or hexyl, methylsulfinyl, ethylsulfinyl, n- or isopropylsulfinyl, n-, iso-, sec- or tert-butylsulfinyl, methylsulfonyl, ethylsulfonyl, n- or isopropylsulfonyl, n-, iso-, sec- or tert-butylsulfonyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, cyclopropyl, cyclopentyl, cyclohexyl, trifluoromethyl, trichloromethyl, trifluoroethyl, difluoromethylthio, difluorochloromethylthio, trifluoromethylthio, trifluoromethylsulfinyl, trifluoromethylsulfonyl, trifluoromethoxymethyl; $-CH_2-CHO$,
30 $-CH_2CH_2-CHO$, $-CH_2-CO-CH_3$, $-CH_2-CO-CH_2CH_3$, $-CH_2-CO-CH(CH_3)_2$,
 $-CH_2CH_2-CO-CH_3$, $-CH_2CH_2-CO-CH_2CH_3$, $-CH_2CH_2-CO-CH(CH_3)_2$,
 $-CH_2-C(O)OCH_3$, $-CH_2-C(O)OCH_2CH_3$, $-CH_2-C(O)OCH(CH_3)_2$,
 $-CH_2CH_2-C(O)OCH_3$, $-CH_2CH_2-C(O)OCH_2CH_3$, $-CH_2CH_2-C(O)OCH(CH_3)_2$,
 $-CH_2-CO-CF_3$, $-CH_2-CO-CCl_3$, $-CH_2-CO-CH_2CF_3$, $-CH_2-CO-CH_2CCl_3$,

$-\text{CH}_2\text{CH}_2-\text{CO}-\text{CH}_2\text{CF}_3$, $-\text{CH}_2\text{CH}_2-\text{CO}-\text{CH}_2\text{CCl}_3$, $-\text{CH}_2-\text{C}(\text{O})\text{OCH}_2\text{CF}_3$,
 $-\text{CH}_2-\text{C}(\text{O})\text{OCF}_2\text{CF}_3$, $-\text{CH}_2-\text{C}(\text{O})\text{OCH}_2\text{CCl}_3$, $-\text{CH}_2-\text{C}(\text{O})\text{OCCl}_2\text{CCl}_3$,
 $-\text{CH}_2\text{CH}_2-\text{C}(\text{O})\text{OCH}_2\text{CF}_3$, $-\text{CH}_2\text{CH}_2-\text{C}(\text{O})\text{OCF}_2\text{CF}_3$, $-\text{CH}_2\text{CH}_2-\text{C}(\text{O})\text{OCH}_2\text{CCl}_3$,
 $-\text{CH}_2\text{CH}_2-\text{C}(\text{O})\text{O}-\text{CCl}_2\text{CCl}_3$; $-\text{COR}^5$, $-\text{CONR}^6\text{R}^7$ or $-\text{CH}_2\text{NR}^8\text{R}^9$,

5 R^5 represents hydrogen, methyl, ethyl, n- or iso-propyl, tert-butyl, methoxy, ethoxy, tert-butoxy, cyclopropyl; trifluoromethyl, trifluoromethoxy; or $-\text{COR}^{10}$,

R^6 and R^7 independently of one another each represent hydrogen, methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, cyclopropyl, cyclopentyl, cyclohexyl; trifluoromethyl, trichloromethyl, trifluoroethyl, trifluoromethoxymethyl,

10

R^6 and R^7 furthermore together with the nitrogen atom to which they are attached, represent a saturated heterocycle selected from the group consisting of morpholine, thiomorpholine and piperazine, which heterocycle may optionally be mono- to tetra-substituted, identically or differently, by fluorine, chlorine, bromine or methyl and where the piperazine additionally at the second nitrogen atom may be substituted by R^{11} ,

15

R^8 and R^9 independently of one another each represent hydrogen, methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, cyclopropyl, cyclopentyl, cyclohexyl; trifluoromethyl, trichloromethyl, trifluoroethyl, trifluoromethoxymethyl,

20


R^8 and R^9 furthermore together with the nitrogen atom to which they are attached, represent a saturated heterocycle selected from the group consisting of morpholine, thiomorpholine and piperazine, which heterocycle may optionally be mono- to tetra-substituted, identically or differently, by fluorine, chlorine, bromine or methyl and where the piperazine additionally at the second nitrogen atom may be substituted by R^{11} ,

25

R^{10} represents hydrogen, methyl, ethyl, n- or iso-propyl, tert-butyl, methoxy, ethoxy, n- or iso-propoxy, tert-butoxy, cyclopropyl; trifluoromethyl, trifluoromethoxy,

R^{11} represents hydrogen, methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl,

A represents a radical of the formula (A1)



(A1), wherein

30

R^{12} represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, ethyl, iso-propyl, monofluoromethyl, monofluoroethyl, difluoromethyl, trifluoromethyl, difluorochloromethyl, trichloromethyl, dichloromethyl, cyclopropyl,


methoxy, ethoxy, trifluoromethoxy, trichloromethoxy, methylthio, ethylthio, trifluoromethylthio or difluoromethylthio and

R^{13} represents hydrogen, fluorine, chlorine, bromine, iodine or methyl and

R^{14} represents hydrogen, methyl, ethyl, iso-propyl, trifluoromethyl, difluoromethyl, hydroxymethyl, hydroxyethyl or phenyl,

or

A represents a radical of the formula (A2)




(A2), wherein

R^{15} and R^{16} independently of one another each represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl, difluoromethyl, trifluoromethyl, difluorochloromethyl or trichloromethyl and

R^{17} represents fluorine, chlorine, bromine, cyano, methyl, trifluoromethyl, trifluoromethoxy, difluoromethoxy, difluorochloromethoxy or trichloromethoxy,

or

A represents a radical of the formula (A4)




(A4), wherein

R^{21} represents hydrogen, fluorine, chlorine, bromine, iodine, hydroxyl, cyano, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, difluoromethyl, trifluoromethyl, difluorochloromethyl, trichloromethyl, trifluoromethoxy, difluoromethoxy, difluorochloromethoxy, trichloromethoxy, trifluoromethylthio, difluoromethylthio, difluorochloromethylthio or trichloromethylthio,

or

A represents a radical of the formula (A5)



(A5), wherein


R^{22} represents fluorine, chlorine, bromine, iodine, hydroxyl, cyano, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, trifluoromethyl, difluoromethyl, difluorochloromethyl, trichloromethyl, methoxy, ethoxy, methylthio, ethylthio, difluoromethylthio, trifluoromethylthio, tri-

fluoromethoxy, difluoromethoxy, difluorochloromethoxy or trichloromethoxy and

R^{23} represents hydrogen, fluorine, chlorine, bromine, iodine, cyano, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, trifluoromethyl, difluoromethyl, difluorochloromethyl, trichloromethyl, methoxy, ethoxy, methylthio, ethylthio, trifluoromethoxy, difluoromethoxy, difluorochloromethoxy, trichloromethoxy, methylsulphinyl or methylsulphonyl,

or

A represents a radical of the formula (A6)



R^{24} represents methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl and


R^{25} represents methyl,

Q^3 represents a sulphur atom or CH_2 ,

P represents O,

or

A represents a radical of the formula (A9)



R^{28} and R^{29} independently of one another each represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl and

R^{30} represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl,

or

A represents a radical of the formula (A11)




R^{34} represents hydrogen, fluorine, chlorine, bromine, amino, methylamino, dimethylamino, cyano, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl and

R^{35} represents fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl,

or

A represents a radical of the formula (A17)



(A17), wherein

R^{43} preferably represents fluorine, chlorine, bromine, iodine, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, trifluoromethyl, difluoromethyl, difluorochloromethyl, trichloromethyl,


excluded compounds of the formula (I), in which

R represents hydrogen and

R^1 , R^2 and R^3 independently of one another each represents hydrogen, fluorine, chlorine, bromine; methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl; or trifluoromethyl or trifluoroethyl; and

R^4 represents hydrogen and

A represents a radical of the formula (A1)



(A1), wherein

R^{12} represents fluorine, chlorine, bromine, iodine, methyl, ethyl, iso-propyl, monofluoromethyl, monofluoroethyl, difluoromethyl, trifluoromethyl, difluorochloromethyl, trichloromethyl, dichloromethyl and

R^{13} represents hydrogen and

R^{14} represents methyl,

or

A represents a radical of the formula (A2)




(A2), wherein

R^{15} and R^{16} independently of one another each represent hydrogen, methyl or ethyl and

R^{17} represents fluorine, chlorine, bromine, methyl, ethyl, or trifluoromethyl,

or

A represents a radical of the formula (A4)




(A4), wherein

R²¹ represents fluorine, chlorine, bromine, iodine, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, difluoromethyl, trifluoromethyl, difluorochloromethyl, trichloromethyl,

or

A represents a radical of the formula (A5)




(A5), wherein

R²² represents fluorine, chlorine, bromine, iodine and

R²³ represents hydrogen,

or

A represents a radical of the formula (A6)



(A6), wherein


R²⁴ represents methyl and

Q³ represents a sulphur or CH₂,

p represents 0,

or

A represents a radical of the formula (A9)




(A9), wherein

R²⁸ and R²⁹ independently of one another each represent hydrogen, methyl or ethyl
and

R³⁰ represents methyl,

or

A represents a radical of the formula (A11)



(A11), wherein

R^{34} represents hydrogen, methyl or ethyl and


R^{35} represents fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.

5 4. Pyridinylanilides of the formula (I) according to Claims 1, 2 or 3, in which R^4 represents hydrogen.

5. Pyridinylanilides of the formula (I) according to Claims 1, 2 or 3, in which R represents hydrogen.

10

6. Pyridinylanilides of the formula (I-12)



(I-12)

in which

R, R^4 and A are as defined in Claims 1, 2 or 3 and

15

R^{1a} represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thio-carbamoyl;

or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;

20

or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;

or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;

25

or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;

or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;


30

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;
or represents the grouping $-C(Q^1)=N-Q^2$, wherein

Q^1 represents hydrogen, hydroxyl or C_1 - C_4 -alkyl, C_1 - C_4 -halogenoalkyl having 1 to 9 identical or different halogen atoms, or C_3 - C_6 -cycloalkyl and

Q^2 represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -alkylamino, di(C_1 - C_4 -alkyl)amino or phenyl; or represents C_2 - C_4 -alkenyloxy or C_2 - C_4 -alkynyloxy, represents phenyl, phenoxy, phenylthio, benzoyl, benzoylthio, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched C_1 - C_4 -alkyl and C_1 - C_4 -alkoxy.

7. Pyridinylanilides of the formula (I-13)



in which

R , R^4 and A are as defined in Claims 1, 2 or 3 and

R^{1a} represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thio-carbamoyl;

or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkyl-sulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;

or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;

or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;

or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;

or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;


or represents the grouping $-C(Q^1)=N-Q^2$, wherein

Q^1 represents hydrogen, hydroxyl or C_1 - C_4 -alkyl, C_1 - C_4 -halogenoalkyl having 1 to 9 identical or different halogen atoms, or C_3 - C_6 -cycloalkyl and

Q^2 represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -alkylamino, di(C_1 - C_4 -alkyl)amino or phenyl; or represents C_2 - C_4 -alkenyloxy or C_2 - C_4 -alkynyloxy,

represents phenyl, phenoxy, phenylthio, benzoyl, benzoylethenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclylalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched C_1 - C_4 -alkyl and C_1 - C_4 -alkoxy.

8. Pyridinylanilides of the formula (I-14)



in which

R , R^4 and A are as defined in Claims 1, 2 or 3 and

R^{1a} and R^{2a} independently of one another each represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl;

or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;

or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;

or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;

or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;

or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;


or represents the grouping $-C(Q^1)=N-Q^2$, wherein

Q^1 represents hydrogen, hydroxyl or C_1 - C_4 -alkyl, C_1 - C_4 -halogenoalkyl having 1 to 9 identical or different halogen atoms, or C_3 - C_6 -cycloalkyl and

Q^2 represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -alkylamino, di(C_1 - C_4 -alkyl)amino or phenyl; or represents C_2 - C_4 -alkenyloxy or C_2 - C_4 -alkynyloxy,

represents phenyl, phenoxy, phenylthio, benzoyl, benzoylethenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclylalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched C_1 - C_4 -alkyl and C_1 - C_4 -alkoxy.

9. Pyridinylanilides of the formula (I-15)




in which

R , R^4 and A are as defined in Claims 1, 2 or 3 and

R^{1a} and R^{2a} independently of one another each represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl;
or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;
or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;
or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;
or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;
or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;
or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;
or represents the grouping $-C(Q^1)=N-Q^2$, wherein
 Q^1 represents hydrogen, hydroxyl or C_1 - C_4 -alkyl, C_1 - C_4 -halogenoalkyl having 1 to 9 identical or different halogen atoms, or C_3 - C_6 -cycloalkyl and
 Q^2 represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -alkylamino, di(C_1 - C_4 -alkyl)amino or phenyl; or represents C_2 - C_4 -alkenyloxy or C_2 - C_4 -alkynyloxy,
represents phenyl, phenoxy, phenylthio, benzoyl, benzoylphenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclylalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched C_1 - C_4 -alkyl and C_1 - C_4 -alkoxy.

10. Pyridinylanilides of the formula (I-16)



in which

R, R⁴ and A are as defined in Claims 1, 2 or 3 and

R^{1a}, R^{2a} and R^{3a} independently of one another each represents halogen, cyano, nitro, amino,

hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl;

or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;

or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;

or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;

or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;

or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;

or represents the grouping -C(Q¹)=N-Q², wherein

Q¹ represents hydrogen, hydroxyl or C₁-C₄-alkyl, C₁-C₄-halogenoalkyl having 1 to 9 identical or different halogen atoms, or C₃-C₆-cycloalkyl and

Q² represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents C₁-C₄-alkyl or C₁-C₄-alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino, di(C₁-C₄-alkyl)amino or phenyl; or represents C₂-C₄-alkenyloxy or C₂-C₄-alkynyloxy,

represents phenyl, phenoxy, phenylthio, benzoyl, benzoylthio, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclylalkyl

having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched C₁-C₄-alkyl and C₁-C₄-alkoxy.

- 5 11. Process for preparing pyridinylanilides of the formula (I) according to Claim 1, characterized in that


- a) carboxylic acid derivatives of the formula (II)



in which

- 10 X¹ represents halogen or hydroxyl and
A is as defined in Claim 1,

are reacted with amines of the formula (III)




15 in which

R, R¹, R², R³ and R⁴ are as defined in Claim 1,

if appropriate in the presence of a catalyst, if appropriate in the presence of a condensing agent, if appropriate in the presence of an acid binder and if appropriate in the presence of a diluent,

- 20 or

- b) halogeno-carboxamides of the formula (IV)




in which

R, R⁴ and A are as defined in Claim 1, and

- 25 X² represents bromine or iodine,

are reacted with boronic acid derivatives of the formula (V)



in which


R^1 , R^2 and R^3 are as defined in Claim 1, and

A^1 and A^2 each represent hydrogen or together represent tetramethylethylene,

in the presence of a catalyst, if appropriate in the presence of an acid binder and if appropriate in the presence of a diluent,

or

c) carboxamide boronic acid derivatives of the formula (VI)




in which

R , R^4 and A are as defined in Claim 1, and

A^3 and A^4 each represent hydrogen or together represent tetramethylethylene,

are reacted with pyridinyl derivatives of the formula (VII)




in which

R^1 , R^2 and R^3 are as defined in Claim 1,

in the presence of a catalyst, if appropriate in the presence of an acid binder and if appropriate in the presence of a diluent,

or

d) halogeno-carboxamides of the formula (IV)




in which

R , R^4 and A are as defined in Claim 1, and

X^2 represents bromine or iodine,

are reacted with pyridinyl derivatives of the formula (VII)




in which

R^1 , R^2 and R^3 are as defined in Claim 1,

in the presence of a palladium or platinum catalyst and in the presence of 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bis-1,3,2-dioxaborolane [bis(pinacolato)diboron], if appropriate in the presence of an acid binder and if appropriate in the presence of a diluent,

or

e) pyridinylanilides of the formula (I-1)



in which

R , R^1 , R^2 , R^3 and A are as defined in Claim 1,

are reacted with halogenides of the formula (VIII)




in which

X^3 represents chlorine, bromine or iodine,

R^{4a} represents C_1 - C_8 -alkyl, C_1 - C_6 -alkylsulfinyl, C_1 - C_6 -alkylsulfonyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_3 - C_8 -cycloalkyl; C_1 - C_6 -halogenoalkyl, C_1 - C_4 -halogenoalkylthio, C_1 - C_4 -halogenoalkylsulfinyl, C_1 - C_4 -halogenoalkylsulfonyl, halogeno- C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_3 - C_8 -halogenocycloalkyl having in each case 1 to 9 fluorine-, chlorine- and/or bromine atoms; formyl- C_1 - C_3 -alkyl, (C_1 - C_3 -alkyl)carbonyl- C_1 - C_3 -alkyl, (C_1 - C_3 -alkoxy)carbonyl- C_1 - C_3 -alkyl; (C_1 - C_3 -halogenoalkyl)carbonyl- C_1 - C_3 -alkyl, (C_1 - C_3 -halogenoalkoxy)carbonyl- C_1 - C_3 -alkyl having in each case 1 to 7 fluorine-, chlorine- and/or bromine atoms,


- (C₁-C₃-alkyl)carbonyl-C₁-C₃-halogenoalkyl, (C₁-C₃-alkoxy)carbonyl-C₁-C₃-halogenoalkyl having in each case 1 to 6 fluorine-, chlorine- and/or bromine atoms, (C₁-C₃-halogenoalkyl)carbonyl-C₁-C₃-halogenoalkyl, (C₁-C₃-halogenoalkoxy)carbonyl-C₁-C₃-halogenoalkyl having in each case 1 to 13 fluorine-, chlorine- and/or bromine atoms; -COR⁵, -CONR⁶R⁷ or -CH₂NR⁸R⁹,
 5 R⁵, R⁶, R⁷, R⁸ and R⁹ are as defined in Claim 1,
 in the presence of a base and in the presence of a diluent.

12. Compositions for controlling unwanted microorganisms, characterized in that they comprise
 10 at least one pyridinylanilide of the formula (I) according to Claim 1, in addition to extenders and/or surfactants.
13. Use of pyridinylanilides of the formula (I) according to Claim 1 for controlling unwanted
 15 microorganisms.
14. Method for controlling unwanted microorganisms, characterized in that pyridinylanilides of
 the formula (I) according to Claim 1 are applied to the microorganisms and/or their habitats.
15. Process for preparing compositions for controlling unwanted microorganisms, characterized
 20 in that pyridinylanilides of the formula (I) according to Claim 1 are mixed with extenders and/or surfactants.
16. Amines of the formula (III)



- 25 in which
 R, R¹, R², R³ and R⁴ are as defined in Claim 1,
 excluded compounds of the formula (III), in which
 R represents hydrogen and ^
 R¹, R² and R³ independently of one another each represents hydrogen, halogen, straight-chain
 30 or branched alkyl having 1 to 4 carbon atoms or straight-chain or branched
 halogenoalkyl having 1 to 4 carbon atoms; and R⁴ represents hydrogen.

17. Amines of the formula




in which

R and R⁴ are as defined in Claims 1, 2 or 3 and

- 5 R^{1a} represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thio-carbamoyl;
 or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkyl-sulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;
 10 or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;
 or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;
 15 or represents in each case straight-chain or branched halogenoalkenyl or halogeno-alkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;
 or represents in each case straight-chain or branched alkylamino, dialkylamino, alkyl-carbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocar-bonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms
 20 in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;
 or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;
 or represents the grouping -C(Q¹)=N-Q², wherein
 25 Q¹ represents hydrogen, hydroxyl or C₁-C₄-alkyl, C₁-C₄-halogenoalkyl having 1 to 9 identical or different halogen atoms, or C₃-C₆-cycloalkyl and
 Q² represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents C₁-C₄-alkyl or C₁-C₄-alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino, di(C₁-C₄-alkyl)amino or phenyl; or represents C₂-C₄-alkenyloxy or C₂-C₄-alkynyloxy,
 30 represents phenyl, phenoxy, phenylthio, benzoyl, benzoylphenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclylalkyl

having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched C₁-C₄-alkyl and C₁-C₄-alkoxy.

5 18. Amines of the formula



in which

R and R⁴ are as defined in Claims 1, 2 or 3 and

R^{1a} represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thio-
10 carbamoyl;

or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkyl-
sulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;

or represents in each case straight-chain or branched alkenyl or alkenyloxy having in
15 each case 2 to 6 carbon atoms;

or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each
case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;

or represents in each case straight-chain or branched halogenoalkenyl or halogeno-
20 alkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;

or represents in each case straight-chain or branched alkylamino, dialkylamino, alkyl-
carbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocar-
bonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms
25 in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;


or represents the grouping -C(Q¹)=N-Q², wherein

Q¹ represents hydrogen, hydroxyl or C₁-C₄-alkyl, C₁-C₄-halogenoalkyl having 1
30 to 9 identical or different halogen atoms, or C₃-C₆-cycloalkyl and

Q² represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents C₁-C₄-alkyl or C₁-C₄-alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino, di(C₁-C₄-alkyl)amino or phenyl; or represents C₂-C₄-alkenyloxy or C₂-C₄-alkynyloxy, represents phenyl, phenoxy, phenylthio, benzoyl, benzoylethenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclylalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched C₁-C₄-alkyl and C₁-C₄-alkoxy.

10

19. Amines of the formula



in which

R and R⁴ are as defined in Claims 1, 2 or 3 and

- 15 R^{1a} and R^{2a} independently of one another each represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl;
or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;
20 or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;
or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;
25 or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;
or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms
- 30

in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;


or represents the grouping $-C(Q^1)=N-Q^2$, wherein

Q^1 represents hydrogen, hydroxyl or C_1 - C_4 -alkyl, C_1 - C_4 -halogenoalkyl having 1 to 9 identical or different halogen atoms, or C_3 - C_6 -cycloalkyl and

Q^2 represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -alkylamino, di(C_1 - C_4 -alkyl)amino or phenyl; or represents C_2 - C_4 -alkenyloxy or C_2 - C_4 -alkynyloxy,

represents phenyl, phenoxy, phenylthio, benzoyl, benzoylphenyl, cinnamoyl, heterocyclyl or phenylalkyl; phenylalkyloxy, phenylalkylthio or heterocyclalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched C_1 - C_4 -alkyl and C_1 - C_4 -alkoxy.

20. Amines of the formula



in which

R and R^4 are as defined in Claims 1, 2 or 3 and

R^{1a} and R^{2a} independently of one another each represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl;

or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;

or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;

or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;

or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;

or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;


or represents the grouping $-C(Q^1)=N-Q^2$, wherein

Q^1 represents hydrogen, hydroxyl or C_1 - C_4 -alkyl, C_1 - C_4 -halogenoalkyl having 1 to 9 identical or different halogen atoms, or C_3 - C_6 -cycloalkyl and

Q^2 represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -alkylamino, di(C_1 - C_4 -alkyl)amino or phenyl; or represents C_2 - C_4 -alkenyloxy or C_2 - C_4 -alkynyloxy,

represents phenyl, phenoxy, phenylthio, benzoyl, benzoylthenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclylalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched C_1 - C_4 -alkyl and C_1 - C_4 -alkoxy.

21. Amines of the formula



in which

R and R^4 are as defined in Claims 1, 2 or 3 and

R^{1a} , R^{2a} and R^{3a} independently of one another each represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxyl, carbamoyl, thiocarbamoyl;

or represents in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulfinyl or alkylsulfonyl having in each case 1 to 8 carbon atoms in the respective alkyl moiety;

or represents in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;

or represents in each case straight-chain or branched halogenoalkyl, halogenoalkoxy, halogenoalkylthio, halogenoalkylsulfinyl or halogenoalkylsulfonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;

or represents in each case straight-chain or branched halogenoalkenyl or halogenoalkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;

or represents in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms in the respective hydrocarbon chain, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chain;

or represents cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;

or represents the grouping $-C(Q^1)=N-Q^2$, wherein

Q^1 represents hydrogen, hydroxyl or C_1 - C_4 -alkyl, C_1 - C_4 -halogenoalkyl having 1 to 9 identical or different halogen atoms, or C_3 - C_6 -cycloalkyl and

Q^2 represents hydroxyl, amino, methylamino, phenyl, benzyl; or represents C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy, each of which is optionally substituted by halogen, cyano, hydroxyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -alkylamino, di(C_1 - C_4 -alkyl)amino or phenyl; or represents C_2 - C_4 -alkenyloxy or C_2 - C_4 -alkynyloxy,

represents phenyl, phenoxy, phenylthio, benzoyl, benzoylphenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclylalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which is optionally mono- to tri-substituted, identically or differently, in the ring moiety by halogen, in each case straight-chain or branched C_1 - C_4 -alkyl and C_1 - C_4 -alkoxy.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.